




EDF Energy

Hunterston B Seaweed Removal

Environmental Appraisal

June 2017



Innovative Thinking - Sustainable Solutions




Page intentionally left blank

Hunterston B Seaweed Removal


Environmental Appraisal

June 2017



Document Information

Document History and Authorisation		
Title	Hunterston B Seaweed Removal	
	Environmental Appraisal	
Commissioned by	EDF Energy	
Issue date	June 2017	
Document ref	R.2827	
Project no	R/4535/1	
Date	Version	Revision Details
14/06/2017	1	Issued for client review
16/06/2017	2	Issued for client use

Prepared (PM)	Approved (QM)	Authorised (PD)
D J Honey	N J Frost	S C Hull
		

Suggested Citation

ABPmer, (2017). Hunterston B Seaweed Removal, Environmental Appraisal, ABPmer Report No. R.2827. A report produced by ABPmer for EDF Energy, June 2017.

Notice

ABP Marine Environmental Research Ltd ("ABPmer") has prepared this document in accordance with the client's instructions, for the client's sole purpose and use. No third party may rely upon this document without the prior and express written agreement of ABPmer. ABPmer does not accept liability to any person other than the client. If the client discloses this document to a third party, it shall make them aware that ABPmer shall not be liable to them in relation to this document. The client shall indemnify ABPmer in the event that ABPmer suffers any loss or damage as a result of the client's failure to comply with this requirement.

Sections of this document may rely on information supplied by or drawn from third party sources. Unless otherwise expressly stated in this document, ABPmer has not independently checked or verified such information. ABPmer does not accept liability for any loss or damage suffered by any person, including the client, as a result of any error or inaccuracy in any third party information or for any conclusions drawn by ABPmer which are based on such information.

All content in this document should be considered provisional and should not be relied upon until a final version marked '*issued for client use*' is issued.

All images on front cover copyright ABPmer apart from seahorse (A J Pearson).

ABPmer

Quayside Suite, Medina Chambers, Town Quay, Southampton, Hampshire SO14 2AQ
T: +44 (0) 2380 711844 W: <http://www.abpmer.co.uk/>

Contents

1	Introduction	1
2	Project Description.....	2
3	Consenting Framework	6
3.1	Marine Licence under the Marine (Scotland) Act 2010.....	6
4	Environmental Receptors.....	8
4.1	Physical Processes.....	8
4.2	Water and Sediment Quality	10
4.3	Nature Conservation and Aquatic Ecology.....	11
4.4	Cumulative and In-Combination Effects	23
5	HRA Screening Assessment.....	24
5.1	Introduction.....	24
5.2	Project Description (Step 2).....	24
5.3	Site Characterisation (Step 3).....	24
5.4	Assessment of Significance (Step 4).....	25
6	Mitigation and Monitoring Plan	26
7	Conclusions	27
8	References	28
9	Abbreviations/Acronyms	30

Appendices

A	Drop-Down Camera Survey Report	32
A.1	Introduction.....	32
A.2	Survey Methodology	32
A.3	Results	34
A.4	References.....	38
A.5	Abbreviations/Acronyms.....	38
B	Water Framework Directive (WFD) Compliance Assessment	68
B.1	Introduction.....	68
B.2	Screening.....	70
B.3	Scoping	76
B.4	Impact Assessment.....	80
B.5	Conclusion	82
B.6	References.....	82
B.7	Abbreviations/Acronyms.....	83

Tables

Table 1.	Co-ordinates of proposed seaweed reduction area	3
Table 2.	Water level data at Millport (northwest of Hunterston)	8
Table 3.	Significant wave height (m) distribution by direction from a location southwest of Little Cumbrae	9
Table 4.	Seaweed biomass within the proposed seaweed reduction area	16
Table 5.	Coastal and marine related nature conservation designated sites within approximately 40 km of the proposed seaweed reduction area	20

Images

Image 1.	Hunterston B cooling water intake jetty	3
Image 2.	Modified nephrops trawl (left) and mobile chain mat (right)	4
Image 3.	Significant wave height (m) distribution by direction from a location southwest of Little Cumbrae (produced using SEASTATES data)	9
Image 4.	Kelp species abundance and local species richness using the SACFOR scale	16

Figures

Figure 1.	Location of Hunterston B Power Station and cooling water intake within the Clyde	2
Figure 2.	Proposed (and 2016) seaweed reduction area and depth contours	4
Figure 3.	Intertidal habitat in the vicinity of the proposed seaweed reduction area	11
Figure 4.	EMODnet habitat map in the vicinity of the proposed seaweed reduction area	13
Figure 5.	Biotopes recorded during drop-down camera survey (May 2017) in the vicinity of the proposed seaweed reduction area	14
Figure 6.	Seaweed abundance within the proposed seaweed reduction area based on SACFOR scale	15
Figure 7.	National nature conservation designations in the vicinity of the proposed seaweed reduction area	21
Figure 8.	International nature conservation designations in the vicinity of the proposed seaweed reduction area	22

1 Introduction

ABPmer has been commissioned by Hunterston B Power Station, owned and operated by EDF Energy, to undertake an environmental appraisal of the potential marine environmental effects of seaweed clearance from the vicinity of its cooling water intake. This information will support a marine licence application for seaweed clearance that will be submitted to Marine Scotland as part of the marine licensing process.

Hunterston B Power Station utilises the sea as a source of cooling water for plant systems. The cooling water enters the station by passing through a coarse screen located at the cooling water intake. On a number of occasions, the station has experienced high levels of seaweed ingress onto the cooling water intake screens. The high levels of impingement have reduced water flow through the screens requiring the station to reduce energy generation. The station undertook a limited programme of seaweed clearance in 2016 from within the vicinity of the cooling water intake with the approval of Marine Scotland. This is considered to have led to a subsequent reduction in seaweed impingement. The station therefore proposes to undertake similar works in 2017.

A teleconference between EDF Hunterston, ABPmer and Marine Scotland on 19 May 2017 confirmed that the works did not require a formal environmental impact assessment (EIA), but that an environmental appraisal would be required to support the marine licence application. In addition, consideration would need to be given to potential impacts on Natura 2000 sites, national nature conservation marine protected areas (MPAs) and achievement of Water Framework Directive (WFD) objectives.

A voluntary environmental appraisal has therefore been prepared to provide the relevant information on potential environmental issues in support of the marine licence application.

The environmental appraisal has been structured as follows:

- Section 2:** **Project Description** presents details of the proposed works;
- Section 3:** **Consenting Framework** outlines the consenting framework against which the proposed works have been assessed;
- Section 4:** **Environmental Appraisal** reviews the potential effects of the proposed works on environmental receptors;
- Section 5:** **Habitats Regulations Appraisal (HRA) Screening Assessment** reviews the potential effects of the proposed works on designated features of European/Ramsar sites;
- Section 6:** **Mitigation and Monitoring Plan** highlights measures to reduce/avoid potential environmental impacts and plans to monitor the environment; and
- Section 7:** **Conclusions** presents a summary of the conclusions of the environmental appraisal.

2 Project Description

Hunterston B Power Station utilises the sea as a source of cooling water for plant systems. The cooling water enters the station by passing through a coarse screen located at the cooling water intake (see Figure 1 and Image 1). The design intent of the coarse screen is to minimise marine debris from entering the cooling water system.




Figure 1. Location of Hunterston B Power Station and cooling water intake within the Clyde

Operational experience has shown that strong winds from the south to southwest coincident with a low tide promotes seaweed from leaving the sea floor and gathering on the coarse screen. If the coarse screen is inundated with large volumes of seaweed, the cooling water flow is reduced which challenges the operability of the plant. To mitigate, a preliminary study was undertaken in 2016 which involved assessing and reducing the seaweed growth local to the cooling water intake (see Figure 2 for 2016 seaweed reduction area). Permission for the activity was obtained from Marine Scotland. The study indicated that by reducing the local seaweed growth the amount of seaweed gathering on the coarse screen following storm conditions was significantly reduced.



Source: EDF Energy

Image 1. Hunterston B cooling water intake jetty

Hunterston B Power Station now wishes to obtain a marine licence to permit further removal of seaweed within a proposed seaweed reduction area (Figure 2), delineated by the coordinates provided in Table 1. The proposed seaweed reduction area covers a reduced spatial extent (0.64 km²) compared to the 2016 seaweed reduction area (1.3 km²), based on the findings of a drop-down camera survey in May 2017 (see Appendix A for details of the survey). The removal technique will consist of using a standard fishing net supplemented with a mobile chain mat were practicable, deployed from a suitable commercial fishing vessel.

Table 1. Co-ordinates of proposed seaweed reduction area

Proposed Seaweed Reduction Area Point (See Figure 2)	Co-ordinates (WGS84)	
	Latitude	Longitude
A	55°43'3794N	04°54'3132W
B	55°43'4454N	04°54'6530W
C	55°42'4260N	04°54'6918W
D	55°42'4380N	04°54'3798W

The fishing net is a standard nephrops trawl 460 meshes in circumference made with 80 mm braided netting. This is mounted on 24 m of standard 250 mm rock hoppers rigged to have minimum bottom contact to reduce the disturbance of the seabed and associated marine life. The net will have an approximate weight in the water of 400 kg. The deployment of the net involves a pair of No. 5 Bison trawl doors producing an opening net of approximately 12 to 15 m. The mobile chain mat is made of 12 m of 16 mm chain mounted in a U-shape onto a 100 mm pipe with 450 mm wheels. The mobile mat is designed to reduce ground contact and disturbance to a minimum (Image 2).

Seaweed collected during the activity will be stored on the vessel and contained within 1 tonne rubble style sacks or an open bunded 1,000 litre pallet tank. The contained seaweed would then be craned ashore and disposed of using an approved method.





Figure 2. Proposed (and 2016) seaweed reduction area and depth contours



Source: EDF Energy

Image 2. Modified nephrops trawl (left) and mobile chain mat (right)

Hunterston B Power Station is seeking a one year marine licence for seaweed removal within the proposed seaweed reduction area, based on three discrete seaweed removal campaigns. These campaigns would be expected to take place in July/August 2017 to remove the current accumulation of seaweed, September/October 2017 in advance of the winter storms and April/May 2018 to provide

post-winter clearance. Subject to weather conditions, the working pattern for each campaign will involve up to approximately ten 12-hour working days.

Based on a drop-down camera survey in May 2017, the main species of seaweed present within the proposed seaweed reduction area is the kelp *Saccharina latissima* (formerly *Laminaria saccharina*). This is also the main type of seaweed that collects on the cooling water intake screens (Hunterston B Power Station, *pers. comm.*). Hunterston B Power Station is seeking a marine licence to remove up to a maximum of 50 tonnes wet weight of seaweed during each campaign from within the proposed seaweed reduction area, primarily comprising *S. latissima*. Therefore, this equates to a maximum of 150 tonnes wet weight of seaweed removed during the licence period.

A visual survey of the proposed seaweed reduction area would be undertaken using a drop-down camera to identify the distribution and density of seaweed before each campaign. The survey completed in May 2017 (see Appendix A) will be used to inform the first campaign. Daily monitoring of the amounts of seaweed entering the cooling water intake will also be undertaken to assess the effectiveness of the proposed works.

3 Consenting Framework

The following sections outline the consenting framework against which the proposed works have been assessed.

3.1 Marine Licence under the Marine (Scotland) Act 2010

Under the Marine (Scotland) Act 2010, Scottish Ministers are responsible for the marine licensing system and enforcement in the Scottish inshore region from 0-12 nautical miles (nm). The licensing regime allows regulation of the deposit and removal of substances and objects in the seas around Scotland. Activities must take place in accordance with licence conditions. The removal of wild seaweed constitutes 'a removal of substances/objects from the seabed' and thus requires a marine licence.

In considering an application for a marine licence, the Marine Scotland Licensing Operations Team (MS-LOT) will, as part of the process, take into account Government policy statements and guidance including the Marine Policy Statement (HM Government, 2011) and Scotland's National Marine Plan (The Scottish Government, 2015). As part of the determination process, Marine Scotland will also take account of and give consideration to the need for the following:

- Environmental Impact Assessment (EIA);
- Habitats Regulations Appraisal (HRA);
- Water Framework Directive (WFD) Assessment; and
- Marine Protected Area (MPA) Assessment.

The following sections summarise each of the above and considers whether they are likely to be required to support the marine licence application for the proposed works.

3.1.1 Environmental Impact Assessment

The Environmental Impact Assessment Directive (2011/92/EU) sets out the procedure that must be followed before approval is granted for a range of plans and projects, defined in Annexes I and II of the Directive. Annex I projects are considered to have significant effects on the environment and EIA is mandatory. The potential for significant effects on the environment as a result of Annex II projects, and thus whether an EIA is required, however, is at the discretion of the Competent Authority, in this case Marine Scotland, having regard to criteria set out in Annex III of the Directive.

Wild seaweed removal is not identified under either Annex I or Annex II of the Directive. Therefore, a formal EIA is not required. Marine Scotland has requested that an environmental appraisal is prepared to support the marine licence application (teleconference on 19 May 2017). The topics to be covered by the environmental appraisal have been indicated to include:

- Project description;
- Physical processes;
- Water and sediment quality;
- Nature conservation and aquatic ecology;
 - Benthic habitats and species;
 - Nature conservation designated sites; and
- Mitigation and monitoring plan.

3.1.2 Habitats Regulations Appraisal (HRA)

Where a project is located close to, or within, an area designated or proposed under the Birds¹ and/or Habitats Directives² (European sites) and/or the Ramsar Convention³ (Ramsar sites), the requirements of the Conservation (Natural Habitats, &c.) Regulations 1994 (the Habitats Regulations) apply. In essence, this requires the lead Competent Authority, in this case Marine Scotland, to determine whether the proposed works are likely to have a significant effect on a European/Ramsar site and, if so, to undertake an Appropriate Assessment (AA) of the implications of the proposals in the light of the site's conservation objectives.

The nearest European/Ramsar site to the scheme is the Renfrewshire Heights Special Protection Area (SPA), located approximately 12 km northeast from Hunterston. However, this site is designated for terrestrial features (Scottish Natural Heritage (SNH), 2007). The nearest marine related site is the Inner Clyde SPA and Ramsar site, located more than 25 km to the northeast of the proposed works, while the Inner Hebrides and the Minches Special Area of Conservation (SAC) is over 40 km to the west. An HRA screening assessment is presented in Section 5.

3.1.3 Water Framework Directive

The Water Framework Directive (WFD) (2000/60/EEC) establishes a framework for the management and protection of Europe's water resources. It is implemented in Scotland through the Water Environment Water Services (Scotland) Act 2003 and the Water Environment (Controlled Activities) (Scotland) Regulations 2011, more commonly known as the Controlled Activity Regulations (CAR). The overall objective of the WFD is to achieve "good ecological and good chemical status" in all inland and coastal waters. The initial deadline to meet this objective was 2015; however, in cases where it was not possible to do so due to disproportionate expense, natural conditions or technical feasibility, the deadline to achieve "good ecological and good chemical status" is extended to 2027.

The proposed works are located within the Largs Channel (Fairlie Roads) coastal water body. It is necessary to consider whether the proposed works might compromise achievement of WFD objectives for this and/or adjacent water bodies. A WFD compliance assessment is presented in Appendix B.

3.1.4 Nature Conservation Marine Protected Areas

The Marine (Scotland) Act 2010 provides for the designation of nature conservation Marine Protected Areas (MPAs) in Scottish inshore waters (within 12 nm of the territorial baseline) to protect features (habitats and species) considered to be of national importance. Thirty nature conservation MPAs were designated in 2014 covering both Scottish inshore and offshore waters. Under Section 83 of the Act, when considering granting a marine licence, Marine Scotland has to take account of potential impacts to MPA features.

The proposed works are over 10 km from the nearest nature conservation MPA. The South Arran MPA, at approximately 16 km to the southwest of Hunterston, is designated due to a diversity of habitats and species including maerl beds, kelp and seagrass (SNH, 2014). The Southannan Sands Site of Special Scientific Interest (SSSI) is located approximately 0.5 km north and east of the proposed works. This is a nationally designated site due to intertidal sandflat habitat (SNH, 2013). Nature conservation designations are discussed in Section 4.3.

¹ Council Directive 79/409/EEC on the Conservation of Wild Birds.

² Council Directive 92/43/EEC on the Conservation of Natural Habitats and Wild Fauna.

³ Ramsar Convention on Wetlands – Conservation on Wetlands of International Importance, especially on Waterfowl Habitat.

4 Environmental Receptors

There is limited potential for environmental effects to arise from the seaweed removal activity at Hunterston B given the scale and nature of the proposed works. This section considers any potential effects in the context of the following receptors:

- Physical processes;
- Water and sediment quality;
- Nature conservation and aquatic ecology;
 - Benthic habitat and species; and
 - Nature conservation designated sites.

4.1 Physical Processes

Hunterston B Power Station is located at the southern end of the Inner Firth of Clyde. Water level data at Hunterston, based on levels at Millport (approximately 3.5 km northwest of Hunterston), is described on the United Kingdom Hydrographic Office (UKHO) Admiralty Chart 'Firth of Clyde Hunterston Channel and Rothesay Sound' (Chart No. 1867). These data are reproduced in Table 2 and show a spring and neap tidal range of 2.9 m and 1.8 m, respectively.

Table 2. Water level data at Millport (northwest of Hunterston)

Tidal Level	Chart Datum (CD) (m)	Ordnance Datum (Newlyn) (ODN) (m)
Mean High Water Springs (MHWS)	3.4	1.78
Mean High Water Neaps (MHWN)	2.8	1.18
Mean Low Water Springs (MLWS)	0.5	-1.12
Mean Low Water Neaps (MLWN)	1.0	-0.62
Mean Sea Level (MSL)	1.9	0.28

Source: UKHO Chart No. 1867

The Inner Firth of Clyde is characterised by relatively weak currents that are moderated by the wind. This area is relatively shallow, with some deeper channels in excess of 50 m. Tidal flows tend to be low, generally below 0.9 knots (~0.46 m/s). However, marginally faster flow rates are observed through restricted channels such as those around the Isle of Bute and Great Cumbrae. Tidal flows at Hunterston are described in the UKHO Chart No. 1867, with two tidal diamonds in close proximity (both in the Hunterston Channel). Tidal diamond A (55°44'6N, 04°54'0W) is situated approximately 2.0 km north, while tidal diamond B (55°43'7N, 04°55'6W) is approximately 1.5 km west of Hunterston and immediately adjacent to the proposed seaweed reduction area. Over a tidal cycle, the tidal excursion in waters off Hunterston is around 10 km, based on the tidal information presented in the UKHO Chart No. 1867.

Tidal diamonds A and B exhibit an approximate north-easterly flow during the flooding tide and an approximate south-westerly flow during the ebbing tide, following the orientation of the Hunterston Channel. For tidal diamond A, peak flow in both spring and neap tides occurs three hours after high water with flow velocities of 1.4 and 0.9 knots (0.72 and 0.46 m/s), respectively. The maximum current velocities of flooding tides occur five hours before high water, with flows of 1.1 knots (0.57 m/s) during spring tides and 0.7 knots (0.36 m/s) during neap tides. Flow speeds for tidal diamond B are notably lower compared to tidal diamond A, with peak flows during spring and neap tides of 0.7 and 0.4 knots (0.36 and 0.21 m/s) occurring five hours after high water.

Table 3. Significant wave height (m) distribution by direction from a location southwest of Little Cumbrae

Significant Wave Height (m)	Directional Sector Proportion (%)								Total
	N	NE	E	SE	S	SW	W	NW	
0 to 1	4	3	6	9	27	12	13	16	89
1 to 2	0	0	0	1	4	2	2	1	10
2 to 3	0	0	0	0	0	0	0	0	1
3 to 4	0	0	0	0	0	0	0	0	0
4 to 5	0	0	0	0	0	0	0	0	0
Total	4	3	6	10	32	14	16	16	100

Source: SEASTATES

SEASTATES⁴ is ABPmer’s metocean information service which contains a hindcast of hourly wave conditions for the period 1979 to 2015 inclusive (ABPmer, 2013a). Data have been extracted in the vicinity of Hunterston, specifically from a location southwest of Little Cumbrae (55°40’4N, 04°56’13W), and analysed to determine the distribution of wave height by direction as shown in Table 3 and Image 3. The data indicates that waves predominantly originate from a southerly direction (32%), with notable contributions from the southwest, west and northwest (46% combined). However, it should be noted that Little Cumbrae provides a degree of shelter from wave action originating from the west and northwest to the coastline adjacent to Hunterston B, largely restricting potential effects to waves from the south and southwest.




Image 3. Significant wave height (m) distribution by direction from a location southwest of Little Cumbrae (produced using SEASTATES data)

⁴ ABPmer SEASTATES: www.seastates.net (Accessed June 2017).

The coastline at Hunterston B Power Station is west facing and, therefore, wave action from the south would impact littoral habitats/sediments through refraction towards the coast while wave action from a southwest direction could directly impact the coastline at Hunterston. It is these wave conditions, combined with low water tidal states, which are considered to push seaweed towards the Hunterston B cooling water intake.

The removal of seaweed, predominantly *S. latissima*, from the proposed seaweed reduction area has the potential to increase wave energy reaching the adjacent shoreline as seaweed such as kelps can attenuate wave activity. This, in turn, could lead to changes in the morphology of the intertidal area as a result of increased wave energy acting on the shoreline. The extent of wave dampening is strongly influenced by the morphology, and drag co-efficient, and density of the dominant kelp species; thus, the magnitude of protection provided varies with species, and therefore may also vary with location (Gaylord et al. 2007).

The proposed seaweed reduction area is located in a relatively sheltered area of the Inner Clyde, although extreme wave heights of greater than 2 m do occasionally occur. The abundance and density of *S. latissima* varies seasonally, with higher biomass present in summer and lower biomass present in winter and early spring as a result of removal by natural storm events in autumn and winter. The wave attenuating effects of *S. latissima* will thus be lower during the winter when more extreme and more frequent storm events will occur. It is thus considered that naturally occurring *S. latissima* will play a relatively limited role in wave attenuation in this area of the Inner Clyde.

The proposed seaweed reduction area does not extend all the way to the shoreline. Following seaweed clearance activity, a strip of seaweed will remain on the shoreward side of the reduction area which will continue to play a role in wave dampening. Much of the shoreline adjacent to the seaweed reduction area comprises rocky shore and boulders/cobbles. Such shorelines will be relatively insensitive to changes in morphology in response to minor changes in wave energy. The impact of seaweed removal on the morphology of these shorelines is assessed as **negligible**. The shoreline adjacent to the northern part of the seaweed reduction area is more mixed, comprising boulders/cobbles, gravel and some sand. While the morphology of this shoreline could be slightly more sensitive to changes in wave energy, given the minor nature of the changes in wave energy reaching this shoreline, the impact of seaweed removal on the morphology of this shoreline is assessed as **minor**.

4.2 Water and Sediment Quality

The proposed works are located within the Largs Channel (Fairlie Roads) coastal water body (ID: 200026), while the Seamill and Ardrossan coastal water body (ID: 200024) is located immediately to the south of the proposed seaweed reduction area. In 2015, both coastal water bodies were classified as achieving good overall status⁵. The WFD compliance assessment, presented in Appendix B, concludes that the proposed works at Hunterston are not likely to have a permanent (i.e. non-temporary) effect on the status of WFD parameters that are significant at water body level. The proposed works are therefore not predicted to cause either deterioration to the current status of the Largs Channel (Fairlie Roads) or Seamill and Ardrossan coastal water bodies, nor prevent these water bodies from achieving future WFD objectives. The proposed seaweed reduction activity could lead to small quantities of sediment being raised into suspension through contact with the seabed, although any disturbance will be temporary and sediment will quickly disperse. Given the nature and small scale of the proposed works, water and sediment quality will not be affected and **no impact pathway has been identified**.

⁵ Scotland water body classification results (2015): <http://www.sepa.org.uk/environment/water/classification/classification-results> (Accessed June 2017).

4.3 Nature Conservation and Aquatic Ecology

4.3.1 Benthic Habitats and Species

A number of reviews of the benthic ecology of the Clyde Sea area have been undertaken (e.g. Connor et al. 1998; Wilding et al. 2005). More local reviews of marine habitats in the vicinity of Hunterston B Power Station have also been prepared by Gardline Environmental Ltd (2007) and Cowie (2007). The main habitats and associated species of interest for the appraisal are the subtidal kelp habitats that will be affected by the proposed works and adjacent intertidal areas that could be exposed to indirect impacts as a result of subtidal seaweed removal.

Intertidal Habitats

There is a variety of intertidal habitats in the vicinity of Hunterston B Power Station, ranging from sandflats in more sheltered areas through to boulder and rocky shores in more exposed areas (see Figure 3).




Figure 3. Intertidal habitat in the vicinity of the proposed seaweed reduction area

Hunterston Sands, which lies approximately 2 km northeast of the Hunterston B cooling water intake, is a sandy shore supporting dense lugworm beds (Cowie, 2007). It comprises the biotopes LGS.S.AP.P and LGS.S.Lan (Connor et al. 1998). DHI Ecological Consultancy (2005) reported dense stands of *Zostera noltii* covering approximately 40 hectares (ha) of Hunterston Sands and Southannan Sands.

The area around Brigurd Point (approximately 1.5 km to the north of the Hunterston B cooling water intake) is more exposed than Hunterston Sands. In the upper to mid eulittoral zone there is a mixed habitat matrix of boulders, cobbles, pebbles and sand. The area is characterised by *Semibalanus balanoides* on the boulders and stones together with small clumps of *Mytilus edulis* (Cowie, 2007). Limpets (*Patella vulgata*) are present on larger boulders together with small fucoids (*Fucus vesiculosus*). At the extreme low water level, the substrate is characterised by boulders, cobbles and sand supporting a diverse seaweed community (*F. serratus*, *F. vesiculosus* and encrusting coralline algae) together with encrusting bryozoans and sponges (Gardline Environmental Ltd, 2007). The area has been classified as biotope ELR.Bpat (Connor et al. 1998).

Stoney Port, approximately 1 km to the north of the Hunterston B cooling water intake, is characterised by a complex mosaic of habitats including boulders, cobbles and sand with a larger area of sand towards extreme low water springs. The mid to lower shore is dominated by dense algal cover including *F. vesiculosus* and *Ascophyllum nodosum* and a variety of littorinids (Cowie, 2007).

The coastline adjacent to the cooling water intake jetty (see Image 1) and southwards towards Portencross Castle is more exposed with upper and mid-eulittoral zones comprising steep red sandstone outcrops and boulders. In the lower eulittoral/sublittoral fringe, algae attached to boulders and cobbles form a mosaic with barnacles, littorinids, encrusting bryozoans, sponges and sepolid worms (Gardline Environmental Ltd, 2007).

Subtidal Habitats

There is limited information on the subtidal habitats in the vicinity of Hunterston B Power Station. The European Marine Observation and Data Network (EMODnet)⁶ habitat map classifies the proposed seaweed reduction area as muddy sand/sandy mud with sediments becoming muddier moving offshore (Figure 4), including the following biotopes:

- A3.1: Atlantic and Mediterranean high energy infralittoral rock;
- A5.33: Infralittoral sandy mud; and
- A5.35: Circalittoral sandy mud.

⁶ European Marine Observation and Data Network (EMODnet): <http://emodnet.eu> (Accessed June 2017).




Figure 4. EMODnet habitat map in the vicinity of the proposed seaweed reduction area

The EMODnet habitat map (Figure 4) is supported by the findings of a drop-down camera survey conducted in May 2017 which covered the subtidal area approximately 2 km to the north and 2 km to the south of the Hunterston B cooling water intake (see Appendix A for full details). The main biotopes recorded during the drop-down survey included (see also Figure 5):

- IR.LIR.K.Lsac.Ldig (A3.3131): *Laminaria saccharina* and *Laminaria digitata* on sheltered sublittoral fringe rock;
- SS.SSa.IMuSa (A5.24): Infralittoral muddy sand;
- SS.SMu.ISaMu (A5.33): Infralittoral sandy mud;
- SS.SMp.KSwS.LsacR (A5.521): *Laminaria saccharina* and red seaweeds on infralittoral sediments; and
- SS.SMp.KSwSS.LsacR.S (A5.5213): *Laminaria saccharina* and filamentous red algae on infralittoral sand.




Figure 5. Biotopes recorded during drop-down camera survey (May 2017) in the vicinity of the proposed seaweed reduction area

Kelp

Kelps tend to occur from extreme mean low water springs (MLWS) down to the limit of photic depth (up to 30 m in clear waters; Smale et al. 2013). Dense areas of kelp (*S. latissima*) were recorded at a number of locations in the proposed seaweed reduction area, particularly shallower areas to the north of the cooling water intake. The relative density and distribution of kelp species recorded during the 2017 drop-down camera survey is presented in Figure 6 using the Marine Nature Conservation Review (MNCR) SACFOR scale⁷:

- Super-abundant (S);
- Abundant (A);
- Common (C);
- Frequent (F);
- Occasional (O);
- Rare (R); and
- Less than rare (L).

⁷ SACFOR scale: <http://jncc.defra.gov.uk/page-2684> (Accessed June 2017).

The main kelp species encountered was *S. latissima*. Occasional *Laminaria digitata* were also present. No kelp was observed below 10 m chart datum (CD), suggesting that light is limiting growth beyond this depth. Furthermore, seaweed abundance to the south of the Hunterston B cooling water intake was notably reduced compared to the area north of the intake. It is possible that this is due to the seaweed removal activity undertaken in 2016.




Figure 6. Seaweed abundance within the proposed seaweed reduction area based on SACFOR scale


Maximum standing crop biomass of over 10 kg/m² wet weight has been recorded for *S. latissima* (Krumhansl and Scheibling, 2011). In order to estimate total seaweed biomass within the proposed seaweed reduction area (0.64 km²), based on the findings of the Ma7 2017 drop-down camera survey, it is assumed that 50% of the site (to the north of the Hunterston B cooling water intake) is abundant to super-abundant (1-3 kg/m² wet weight) and 50% of the site (to the south of the Hunterston B cooling water intake) is occasional to frequent (0.2-0.5 kg/m² wet weight). Using these values, it is estimated that total seaweed biomass within the proposed seaweed reduction area ranges between 384 and 1,120 tonnes wet weight (Table 4).

Table 4. Seaweed biomass within the proposed seaweed reduction area

SACFOR Scale	Proportion of Area (%)	Area (km ²)	Seaweed Biomass	
			(kg/m ²)	(Tonnes)
Super-abundant (S) – Abundant (A)	50	0.32	1 – 3	320 – 960
Frequent (F) – Occasional (O)	50	0.32	0.2 – 0.5	64 – 160
Total	100	0.64	-	384 – 1,120

Within the wider Clyde Sea, Wilding et al. (2005) noted that *S. latissima* was widespread with records of extensive beds around Arran and Cumbrae as well as in many of the sea lochs within the Clyde system. *Saccharina latissima* is also widespread throughout Scottish waters (Marine Life Information Network (MarLIN) website)⁸.

Kelps such as *S. latissima* play an important role in coastal marine ecosystems. In particular, they contribute high levels of primary production. For example, Graham et al. (2016) indicated that primary production within *S. latissima* beds can exceed 1,000 g C m²/year, with much of this production entering the carbon cycle as detritus or dissolved organic matter. Kelp species also support notable biodiversity, with higher levels of biodiversity observed in kelp beds than equivalent comparable areas (see Image 4; Smale et al. 2013).



Source: Smale et al. (2013)

Image 4. Kelp species abundance and local species richness using the SACFOR scale

⁸ Sugar kelp (*Saccharina latissima*): <http://www.marlin.ac.uk/species/detail/1375> (Accessed June 2017).

Kelp beds have also been identified as providing important nursery and feeding functions for certain fish and shellfish species. For example, Smale et al. (2013) indicated that kelp forest habitats are vital for the European lobster (*Homarus gammarus*) where it preys on a variety of molluscs and crustaceans. May (2015) identified kelp forests as important for juvenile lobster and juvenile crab (*Cancer pagurus*) around the Isle of Man. Kelp forests also serve as a nursery for many fish species, including Atlantic Cod (*Gadus morhua*) and pollack (*Pollachius pollachius*) (Smale et al. 2013). They are also feeding grounds for fish species such as ballan wrasse (*Labrus bergylta*) and Goldsinny wrasse (*Ctenolabrus rupestris*), which prey on kelp associated invertebrates (Norderhaug et al. 2005), as well as attracting commercially important species such as European sea bass (*Dicentrarchus labrax*), pollack (*P. pollachius*) and conger eels (*Conger conger*) (Smale et al. 2013). In turn, elevated fish densities in kelp forests attract large piscivores, such as large fish, seals and otters.

Maerl

There are no known maerl beds in the vicinity of Hunterston B Power Station (Cowie, 2007) and none were detected during the drop-down camera survey in May 2017 (see Appendix A). The nearest maerl beds are understood to be located in the area of the Tan between Great and Little Cumbrae (Gardline Environmental Ltd, 2007).

Assessment of Potential Impacts

The following impacts to benthic habitats and species have been assessed:

- Direct loss of/damage to kelp habitat within the proposed seaweed reduction area;
- Direct damage to other habitats within the proposed seaweed reduction area;
- Indirect damage to intertidal habitats as a result of changes in wave activity;
- Indirect damage to wider marine ecosystem as a result of removal of primary production;
- Introduction of non-native species; and
- Impact of kelp removal on juvenile and adult fish and shellfish, including commercially important species.

Direct loss of/damage to kelp habitat within the proposed seaweed reduction area

The activity will remove kelp from an area of up to 0.64 km², targeted towards those areas of densest kelp cover based on the findings of a drop-down camera survey in May 2017. The activity will remove some but not all of the kelp from the areas within which the activity occurs. Immediately following the removal, the habitat will support less kelp/less dense kelp than prior to the activity, but the habitat type will not change. Rather some of the ecological functioning of the area could be reduced. Smale et al. (2013) noted that the biodiversity of *S. latissima* beds did not change significantly across the SACFOR scale (see Image 4), suggesting that some aspects of ecological functioning may not be significantly impaired.

Saccharina latissima is likely to regrow rapidly in the targeted areas. This regrowth may offset some of the lost function of the affected habitat. Given the relatively small area over which activity may occur in relation to the overall scale of the resource within the Clyde Sea, the limited impact on ecological structure/functioning and the scope for rapid recovery, the impact on kelp habitat is therefore assessed as **minor**.

Direct damage to other habitats within the proposed seaweed reduction area

No maerl has been identified within the proposed seaweed reduction area based on the May 2017 drop-down camera survey (see also Appendix A). The main biotopes present within the proposed

seaweed reduction area, other than kelp habitat, comprise infralittoral muddy sand (A5.24) and infralittoral sandy mud (A5.33). The removal of *S. latissima* using an otter trawl and/or chain mat will cause some minor disturbance to surface seabed sediments as a result of the physical interaction between the gears and the seabed. The macrofauna and near-surface infauna of subtidal stable muddy sands, sandy muds and muds are susceptible to physical disturbance from bottom fishing gears (i.e. beam trawls, scallop dredges, otter trawls, seine netting, hydraulic suction dredges) (Hall et al. 2008; Kaiser et al. 2002; 2006; Johnson, 2002; Thrush and Dayton, 2002). In general, use of fishing gears that penetrate the substratum to a greater extent (i.e. beam trawls, scallop dredges and demersal trawls) will potentially damage these habitats to a greater degree than activities using lighter gear (i.e. light demersal trawls and seines) (Hall et al. 2008). However, the gears to be used within the proposed seaweed reduction area have been designed to avoid penetration into the seabed, although the gears (particularly the otter trawl doors) could create surface tracks on the softer sedimentary habitats. It is noted that the gears will be less damaging to the sea bed than standard otter trawls or scallop dredges (ABPmer, 2013b).

The abiotic component of sandy mud and muddy sand habitats is considered to have 'High' resistance to the surface abrasion as this pressure is unlikely to alter the habitat type, although there may be some surficial sediment disturbance and the displacement of stones. Recovery is considered to be 'Very High' and the habitat feature is therefore considered to be 'Not Sensitive' to a single event that leads to surface abrasion. Surface disturbance may alter the surface topography of these habitat types, re-suspend sediment and alter sediment characteristics. However, resistance to this pressure is assessed as 'Medium' as the habitat still remains and alterations are confined to surficial layers. In general, any tracks or pits resulting from surface damage would be infilled within six months or sooner by natural hydrodynamic processes and recovery is therefore judged to be 'Very High'. The sensitivity of the abiotic habitat is therefore categorised as 'Low' (ABPmer, 2013b).

Shallow disturbance may lead to injury and mortality of characterising species. Biological recovery is linked to the recovery of the abiotic habitat, which is likely to be rapid in areas where sediments are relatively mobile and will be aided by water transport or active migration of adults. Burrowing polychaetes are considered to be relatively protected from shallow disturbance while species exposed in surface sediments were considered to have lower resistance. Bivalve characterising species are considered to have 'Medium' resistance to this pressure. The high recovery rates of species that characterise these biotopes mean that overall sensitivity was considered to be 'Not Sensitive-Low' (ABPmer, 2013b).

Given the relatively small area affected by the removal activity and the low sensitivity of habitats and associated features to seabed abrasion, the impact on infralittoral muddy sand and infralittoral sandy mud habitats is assessed as negligible.

Indirect damage to intertidal habitats as a result of changes in wave activity

As described in Section 4.1, the largest waves within the study area are from the south and southwest with very small waves from other directions (less than 1 m). Removal of kelp from the shallow sublittoral zone could increase wave propagation onto adjacent shorelines. However, the shoreline adjacent to the proposed seaweed reduction area comprises sandstone outcrops, boulders and cobbles together with patches of sand at Stoney Port. These shorelines will be relatively insensitive to changes in wave propagation. Furthermore, the abundance of *S. latissima* is seasonal with peak biomass in summer and lowest biomass in winter (when the frequency and magnitude of storms and wave action are likely to be greatest). On this basis, the impact of kelp removal on the morphology and ecology of the shoreline adjacent to the proposed seaweed reduction area is assessed as negligible (for those areas of harder shoreline) to minor (for sandy areas of shoreline).

Indirect damage to wider marine ecosystem as a result of removal of primary production

Between June 2015 (following a site event due to seaweed) and December 2015, 231 tonnes of material was collected and subsequently removed from the trash baskets of the Hunterston B cooling water intake (Hunterston B Power Station, *pers. comm.*). The trash baskets are large meshed containers that collect material such as seaweed that has passed through the coarse screens (it should be noted that this figure also includes jellyfish). Up to 150 tonnes wet weight will be removed from within the proposed seaweed reduction area over the one year marine licence (three campaigns at 50 tonnes wet weight each). Assuming approximately 89% of the wet weight is water and 28% of the resulting dry weight is carbon (Gevaert et al. 2001), the activity represents a removal of approximately 4.6 tonnes of carbon from the Clyde system.

Assuming that the volume of kelp removed from the cooling water intake screens reduces as a result of the activity, the overall amount of carbon removed from the Clyde Sea may not increase substantially from the current baseline. Given the small area proposed, this removal of carbon is considered to be **negligible** in the context of the Clyde Sea as a whole.

Introduction of non-native species

The removal of kelp within the proposed seaweed reduction area may create colonising space for the recruitment of non-native species. The proposed seaweed reduction area is 0.64 km² and is naturally disturbed by wave activity during storms on a regular basis. The additional disturbance caused by seaweed removal is considered to be minor. The activity will primarily remove kelp and will cause very little impact to the seabed. On this basis, the increased risk of colonisation by non-native species as a result of the seaweed removal activity is assessed to be very low and the impact is assessed as **negligible**.

The use of a vessel to remove seaweed could pose a risk of introduction of non-native species through discharge of ballast water or biofouling of the vessel's hull. Hunterston B Power Station is committed to using a local fishing vessel to undertake the removal activity. The risk of transfer of non-native species is therefore assessed as **negligible**.

Impact of kelp removal on juvenile and adult fish and shellfish

It is likely that the kelp beds within the proposed seaweed reduction area are of functional value to juvenile cod and pollack and to juvenile lobster and crab. They may also be important feeding areas for adult lobster and a range of fish species. Therefore, removal of kelp within the proposed seaweed reduction area could reduce the functional value of the area for juvenile and adult fish and shellfish. The spatial extent of the proposed seaweed reduction area is small relative to the overall extent of kelp resources in the Clyde Sea. The proposed activity will remove some but not all of the kelp and some residual function as a nursery/feeding area will remain following the activity, while some regrowth of kelp will occur between each campaign. In addition, the proposed seaweed reduction area does not extend all the way to the shoreline. A strip of seaweed will remain on the shoreward side of the reduction area which will also continue to function as a nursery/feeding area.

Given the relatively small area over which seaweed removal may occur in relation to the overall scale of the resource within the Clyde Sea, the limited impact on ecological structure/functioning within proposed seaweed reduction area and the scope for rapid recovery, the impact on fish nursery and feeding function is assessed as **minor**.

4.3.2 Nature Conservation Designated Sites

This section reviews the effects of the proposed works on nature conservation receptors, i.e. designated habitats and species. The proposed works do not directly overlap with any nationally (Figure 7) or internationally (Figure 8) designated nature conservation sites. Table 5 presents a summary of all coastal and marine related designated sites within approximately 40 km of the proposed works and outlines their respective qualifying features.

Table 5. Coastal and marine related nature conservation designated sites within approximately 40 km of the proposed seaweed reduction area

Site Name	Features*	Approx. Distance from Proposed Seaweed Reduction Area (km)
Southannan Sands SSSI	Sandflats (Intertidal marine habitats and saline lagoons: Sandflats)	0.5
Kames Bay SSSI	Sandflats (Habitat: Coastland)	3
Ballochmartin Bay SSSI	Sandflats (Habitat: Coastland)	5
South Arran MPA	Burrowed mud; Kelp and seaweed communities on sublittoral sediment; Maerl beds; Maerl or coarse shell gravel with burrowing sea cucumbers; Seagrass beds; Shallow tide-swept coarse sands with burrowing bivalves; Ocean quahog aggregations	16
Inner Clyde SPA and Ramsar	Non-breeding birds: Redshank (<i>Tringa totanus</i>)	28
Inner Clyde SSSI	Saltmarsh; Non-breeding birds: Cormorant (<i>Phalacrocorax carbo</i>); Eider (<i>Somateria mollissima</i>); Goldeneye (<i>Bucephala clangula</i>); Oystercatcher (<i>Haematopus ostralegus</i>); Red-breasted merganser (<i>Mergus serrator</i>); Red-throated diver (<i>Gavia stellata</i>); Redshank (<i>Tringa totanus</i>)	28
Sound of Gigha dSPA	Non-breeding birds: Eider (<i>Somateria mollissima</i>), Great northern diver (<i>Gavia immer</i>), Red-breasted merganser (<i>Mergus serrator</i>)	37
Inner Hebrides and the Minches SAC	Harbour porpoise (<i>Phocoena phocoena</i>)	42

* Features based on information provided by SNH Sitelink: <http://gateway.snh.gov.uk/sitelink/index.jsp> (Accessed June 2017).

The nearest designated area to the proposed seaweed reduction area is the Southannan Sands SSSI, located approximately 0.5 km to the north of the proposed seaweed reduction area. This site is designated for intertidal sandflat and nationally rare dwarf eelgrass (*Zostera noltei*). The objectives for site management are 'to maintain the extent of the intertidal sandflat habitat by ensuring protection from damaging impacts, in particular any future coast development'. The main impacts are considered to be associated with direct habitat loss and interference with physical processes (SNH, 2013).




Figure 7. National nature conservation designations in the vicinity of the proposed seaweed reduction area

This assessment considers the potential effects on physical processes, water and sediment quality and benthic habitats and species which have been previously assessed for the proposed works (Section 4.1 to 4.3). Specifically, the potential for effects on intertidal areas is considered to be limited to changes in wave activity as detailed in Section 4.3.1. All effects have been assessed as negligible or minor and consequently no significant pathways for effects on Southannan Sands SSSI are identified. No effects are therefore anticipated on Southannan Sands SSSI intertidal features, or the other intertidal SSSIs in the wider area (e.g. Kames Bay and Ballochmartin Bay). Considering the small scale seaweed removal proposed, any effects from the proposed works are likely to be highly localised. The nearest marine related European/Ramsar Site is the Inner Clyde SPA and Ramsar located more than 25 km to the northeast of the proposed works. European and Ramsar sites are considered further within the HRA screening provided in Section 5. Overall, due to the distance of all designated sites from the proposed works, no significant pathways for direct or indirect effects are identified and consequently impacts on nature conservation are assessed as **negligible** or at worst **minor** and not significant.

Under the UK Biodiversity Action Plan (UK BAP), a range of priority habitats were identified as being the most threatened and requiring conservation action. Marine UK BAP priority habitats identified in the vicinity of the proposed seaweed reduction area include coastal saltmarsh, seagrass beds, subtidal

sands and gravels, maerl beds and intertidal mudflats⁹. Local Biodiversity Action Plans (LBAPs) aim to conserve biodiversity through local partnerships, taking into account both national and local priorities. The Ayrshire LBAP includes the marine environment adjacent to Hunterston B Power Station, with a primary objective to maintain and enhance the range of sublittoral sand and gravel communities off the Ayrshire coast¹⁰. As a result of devolution, and new country-level and international drivers and requirements, much of the work previously carried out by the UK BAP is now focussed at a country-level rather than a UK-level. As required under Section 2(4) of the Nature Conservation (Scotland) Act 2004, Scottish Priority Marine Features (PMFs) in the vicinity of the proposed seaweed reduction area include kelp and seaweed communities on subtidal sediment, kelp beds and seagrass beds. There will be no effect on any of the aforementioned habitat types, with the exception of kelp which has been assessed as **minor** in Section 4.3.1.




Figure 8. International nature conservation designations in the vicinity of the proposed seaweed reduction area

⁹ UK Biodiversity Action Plan (BAP) priority habitats: <http://jncc.defra.gov.uk/page-5706> (Accessed June 2017).

¹⁰ Ayrshire Local Biodiversity Action Plan (LBAP): <https://www.south-ayrshire.gov.uk/documents/lbap1.pdf> (Accessed June 2017).

4.4 Cumulative and In-Combination Effects

There are no projects or plans known to be taking place in the vicinity of the proposed seaweed reduction area. In particular, there are no known seaweed harvesting proposals in the Clyde, with the nearest suitable areas located on the west coast of Mull of Kintyre. The potential effects that can be attributed to the proposed works are small scale, localised and considered to be minor at worst for receptors (see Sections 4.1 to 4.3). Therefore, the proposed works are **not predicted to result in any significant adverse cumulative/in-combination effects**.

5 HRA Screening Assessment

5.1 Introduction

This section represents a Habitats Regulations Appraisal (HRA) screening assessment for the proposed works. HRA screening is a requirement under The Conservation of Habitats and Species Regulations 2010 (SI 2010 No. 490) (the 'Habitats Regulations') to determine whether European designated sites are likely to be significantly affected by the proposed works either alone or in combination with other plans and projects. If significant effects are likely, there will be a need for an Appropriate Assessment (AA) to be carried out. The screening, any AA and any subsequent assessment is known as the HRA process. European Commission guidance sets out an approach to undertaking AA as part of a four-stage process (Stages A – D; Oxford Brookes, 2001):

- **Stage A:** Screening to determine the need for an AA;
- **Stage B:** AA and the 'integrity test';
- **Stage C:** Assessment of alternative solutions; and
- **Stage D:** Assessment where adverse effects remain.

The screening assessment (Stage A) is based on Likely Significant Effects (LSE), alone or in combination with other plans or projects. If there are no LSE identified for all the European designated sites, it is considered Stages B, C and D will not be required.

This screening assessment is based on the following four steps (Oxford Brookes, 2001):

- **Step 1:** Determining whether the project or plan is directly connected with or necessary for the management of the site;
- **Step 2:** Describing the project (or plan);
- **Step 3:** Identifying the potential effects on European sites (site characterisation); and
- **Step 4:** Assessing the significance of any effects on European sites.

The proposed works is not directly connected with or necessary for the conservation management of any European sites (Step 1). It is therefore appropriate to proceed to Step 2 of the screening process.

5.2 Project Description (Step 2)

Information on the need for the proposed works and project description are provided within Section 2. This assessment considers that the only viable alternative to the proposed works is the 'do nothing' scenario. However, the proposed works are designed to mitigate the risk of extreme seaweed ingress at the Hunterston B cooling water intake. As identified in Section 4.4, no other plans or project have been identified that would have a significant effect and therefore need to be considered further within this assessment.

5.3 Site Characterisation (Step 3)

The importance of the area is recognised through a number of statutory designations in the wider area. This assessment considers the potential wide ranging foraging area of designated features. The following marine or coastal European/Ramsar designated sites are therefore considered that are located within approximately 40 km of the proposed works:

- Sound of Gigha draft SPA (dSPA);
- Inner Clyde Ramsar site;
- Inner Clyde SPA; and
- Inner Hebrides and the Minches SAC.

As detailed in Section 4.3.2, the site does not overlap with any designated sites and therefore potential effects are associated with indirect effects only. Other sites such as the Renfrewshire Heights and Arran Moors SPAs are also in the wider area but are purely terrestrial sites and have therefore not been considered within this assessment. Details of the existing supporting features of the marine or coastal designated sites are outlined in Table 5. All the European/Ramsar sites identified within 40 km are designated due to birds and harbour porpoise. The primary effects of the proposed works are therefore considered to be associated with loss and/or damage to foraging habitat and species. Section 6 also details mitigation and monitoring plans that are incorporated into the proposed works to minimise effects.

5.4 Assessment of Significance (Step 4)

An assessment of potential effects on physical processes, water and sediment quality and benthic habitats and species has been undertaken in Sections 4.1 to 4.3. All effects are assessed as negligible. Considering the small scale seaweed removal proposed, any effects are likely to be highly localised and no significant effects have been identified. No pathways are therefore identified that have the potential to indirectly effect birds or harbour porpoise within European/Ramsar sites. The proposed works are therefore not considered to have the potential to effects birds associated with The Sound of Gigha dSPA or the Inner Clyde Ramsar site and SPA. Similarly, there would be no indirect effects on harbour porpoise associated with the Inner Hebrides and the Minches SAC.

Overall, considering the proximity of the designated sites from the proposed seaweed reduction area, no pathways are identified that could change the current baseline of European/Ramsar sites in the wider area. The nearest European/Ramsar sites are located over 20 km from Hunterston and any effects associated with the proposed works will be highly localised. The proposed works are therefore not considered to have the potential to effect the integrity of any site conservation objectives. In light of the assessment presented, any effects - whether direct or indirect, permanent or temporary - to interest features are considered to be **negligible** and there is **not considered to be a LSE** on any habitat features or species features associated with the designated sites.

6 Mitigation and Monitoring Plan

To review the distribution and density of seaweed within the proposed seaweed reduction area, a **drop-down camera survey** will be conducted before each campaign. The results of the survey will be reviewed by EDF Energy to ensure the activity is focused on appropriate sections within the proposed seaweed reduction area to remove target species. The methodology would follow the same procedure employed to characterise the proposed seaweed reduction area in May 2017 (see Appendix A), thus also supporting longer-term monitoring of the site.

All seaweed removed from within the proposed seaweed reduction area will be **disposed to land**, as opposed to disposal at sea. This mitigation measure avoids potential adverse impact pathways between the disposed material and other marine habitats. Furthermore, as highlighted in Section 4.3.1, the removal of 150 tonnes of seaweed (wet weight) represents a removal of approximately 4.6 tonnes of carbon from the Clyde system. Assuming that the volume of kelp removed from the cooling water intake screens reduces as a result of the activity, the overall amount of carbon removed from the Clyde Sea may not increase substantially from the current baseline, considered to be negligible in the context of the Clyde Sea as a whole.

The ultimate aim of the proposed works is to mitigate the potential extreme ingress of seaweed at the Hunterston B cooling water intake. Therefore, suitable monitoring of the **amounts of seaweed entering the cooling water intake** will also be undertaken to assess the effectiveness of the activity with a view to reducing the seaweed reduction area over time.

7 Conclusions

Hunterston B Power Station is seeking a one year marine licence to remove up to a maximum of 150 tonnes (wet weight) of seaweed, primarily comprising *Saccharina latissima*, from within a proposed seaweed reduction area in the vicinity of its cooling water intake. The proposed works would involve three separate campaigns during the licence period, each removing up to a maximum of 50 tonnes wet weight of seaweed.

The potential effects that can be attributed to the proposed works are localised and, with the application of mitigation measures, are considered to be minor, negligible or insignificant for all receptors both alone and cumulatively/in-combination with other projects. A summary of how the scheme is compliant with the respective legislative requirements is provided below:

- **Environmental Impact Assessment/Appraisal:** The scale of the proposed works is such that an EIA is not considered necessary. This was agreed with Marine Scotland during a teleconference on 19 May 2017. This is supported by the review of potential impact pathways provided in Section 4 of this environmental appraisal;
- **Habitats Regulations Appraisal:** The proposed works will not have a LSE on any features that are identified within any European/Ramsar sites, either individually or in combination with other plans or projects. An HRA screening assessment is presented in Section 5; and
- **Water Framework Directive:** The scale and nature of the proposed works are very small in scale and unlikely to cause a deterioration or failure of the Largs Channel (Fairlie Roads) or Seamill and Ardrossan coastal water bodies to meet future WFD objectives. A WFD compliance assessment has been prepared in order to comply with the requirements of the WFD and is provided in Appendix B.

8 References

ABPmer. 2013a. SEASTATES Wave Hindcast Model: Calibration and Validation Report. ABP Marine Environmental Research Ltd, Report No. R.2145.

ABPmer. 2013b. Tools for appropriate assessment of fishing and aquaculture activities in marine and coastal Natura 2000 sites. Report III: Intertidal and subtidal muddy sands and sandy muds. Report R.2071. October 2013. Report prepared for the Marine Institute.

Connor, D.W. and Little, M. 1998. Clyde Sea (MNCR Sector 12). In: Marine Nature Conservation Review. Benthic marine ecosystems of Great Britain and the north-east Atlantic. Edited by Hiscock, K. 339-353. Peterborough, Joint Nature Conservation Committee (Coasts and seas of the United Kingdom, MNCR series).

Cowie, P.R., 2007. A synopsis report of the marine biological work carried out in the vicinity of Hunterston power station and potential habitat sensitivities to thermal effluents within a 2 – 5 km radius. A report prepared for British Energy. University of London Marine Biological Station Millport, Isle of Cumbrae, Scotland.

DHI Ecological Consultancy, 2005. Nationally scarce plant survey and woodland national vegetation classification survey of Portencross SSSI, North Ayrshire. Scottish Natural Heritage commissioned Report No. 080.

Gardline Environmental Ltd. 2007. Thermal Plume and Habitat Sensitivity Study Hunterston Nuclear Power Station. British Energy reference 40217233/KMR, December 2007.

Gaylord, B., Rosman, J.H., Reed, D.C., Koseff, J.R., Fram, J., MacIntyre, S., Arkema, K., McDonald, C., Brzezinski, M.A., Largier, J.L. and Monismith, S.G. 2007. Spatial patterns of flow and their modification within and around a giant kelp forest. *Limnology and Oceanography* 52(5): 1838-1852.

Gevaert, F., Davoult, D., Creach, A., Kling, R., Janquin, M., Seuront, L., and Lemoine, Y. 2001. Carbon and nitrogen content of *Laminaria saccharina* in the eastern English Channel: Biometrics and seasonal variations. *Journal of the Marine Biological Association of the United Kingdom* 81(5): 727-734.

Graham, M.H., Fox, M.D. and Hamilton, S.L. 2016. Macrophyte productivity and the provisioning of energy and habitat to nearshore systems. In: *Marine Macrophytes as Foundation Species*, Chapter 7 (pp. 133-152). CRC Press.

Hall, K., Paramour, O.A.L., Robinson, L.A., Winrow-Giffin, A., Frid, C.L.J., Eno, N.C., Dernie, K.M., Sharp, R.A.M., Wyn, G.C. and Ramsay, K. 2008. Mapping the sensitivity of benthic habitats to fishing in Welsh waters – development of a protocol. CCW [Policy Research] Report No: 8/12, 85pp.

HM Government. 2011. UK Marine Policy Statement. HM Government, Northern Ireland Executive, Scottish Government, Welsh Assembly Government. March 2011 London: The Stationery Office. Available online at: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/69322/pb3654-marine-policy-statement-110316.pdf (Accessed June 2017).

Johnson, K.A. 2002. A review of national and international literature on the effects of fishing on benthic habitats. NOAA Technical Memorandum NMFS-F/SPO-57, 72.

- Kaiser, M.J., Clarke, K.R., Hinz, H., Austen, M.C.V., Somerfield, P.J. and Karakassis, I. 2006. Global analysis of response and recovery of benthic biota to fishing. *Marine Ecology Progress Series* 311: 1-14.
- Kaiser, M.J., Collie, J.S., Hall, S.J., Jennings, S. and Poiner, I.R. 2002. Modification of marine habitats by trawling activities: prognosis and solutions. *Fish and Fisheries* 3(2): 114-136.
- Krumhansl K.A. and Scheibling, R.E. 2011. Detrital production in Nova Scotian kelp beds: patterns and processes. *Marine Ecology Progress Series* 421: 67-82.
- May, L. 2015. Identifying habitat associations of European lobster, *Homarus gammarus* (L.) and brown crab, *Cancer pagurus* (L.) in an Isle of Man Marine Protected Area. MSc Thesis, University of Bangor.
- Norderhaug, K.M., Christensen, J.D., Fossa, J.H. and Fredriksen, S. 2005. Fish-macrofauna interactions in a kelp (*Laminaria hyperborea*) forest. *Journal of the Marine Biological Association of the United Kingdom* 85: 1279-1286.
- Oxford Brookes University. 2001. Assessment of plans and projects significantly affecting Natura 2000 sites: Methodological guidance on the provisions of Article 6(3) and 6(4) of the 'Habitats' Directive 92/43/EEC. European Commission DG Environment.
- Scottish Natural Heritage (SNH), 2007. Renfrewshire Heights Special Protection Area (SPA) citation. Available online at: https://gateway.snh.gov.uk/sitelink/siteinfo.jsp?pa_code=8667 (Accessed June 2017).
- Scottish Natural Heritage (SNH), 2013. Southannan Sands Site of Special Scientific Interest (SSSI) citation. Available online at: https://gateway.snh.gov.uk/sitelink/siteinfo.jsp?pa_code=10261 (Accessed June 2017).
- Scottish Natural Heritage (SNH). 2014. South Arran Marine Conservation Order. Available online at: https://gateway.snh.gov.uk/sitelink/siteinfo.jsp?pa_code=10423 (Accessed June 2017).
- Smale, D.A., Burrows, M.T., Moore, P., O'Connor, N. and Hawkins, S.J. 2013. Threats and knowledge gaps provided by kelp forests: a northeast Atlantic perspective. *Ecology and Evolution* 3: 4016-4038.
- The Scottish Government. 2015. Scotland's National Marine Plan. A Single Framework for Managing Our Seas. Marine Scotland. The Scottish Government, Edinburgh 2015. Available online at: <http://www.gov.scot/Resource/0047/00475466.pdf> (Accessed June 2017).
- Trush, S.F. and Dayton, P.K. 2002. Disturbance to marine benthic habitats by trawling and dredging: Implications for Marine Biodiversity. *Annual review of ecology and systematics* 33(1): 449-473.
- Wilding, T.A., Duncan, J., Nickell, L.A., Hughes, D.J., Gontarek, S., Black, K.D. and Sayer, M.D.J. 2005. Synthesis of Information on the Benthos of SEA 6 Clyde Sea Area. Report to the Department of Trade and Industry, Scottish Association for Marine Science, Oban, Scotland, PA37 1QA.





9 Abbreviations/Acronyms

AA	Appropriate Assessment
BAP	Biodiversity Action Plan
CAR	Water Environment (Controlled Activities) (Scotland) Regulations 2011
CD	Chart Datum
dSPA	Draft Special Protection Area
EIA	Environmental Impact Assessment
EMODnet	European Marine Observation and Data Network
HRA	Habitats Regulations Appraisal
LBAP	Local Biodiversity Action Plan
LSE	Likely Significant Effect
MarLIN	Marine Life Information Network
MHWN	Mean High Water Neaps
MHWS	Mean High Water Springs
MLWN	Mean Low Water Neaps
MLWS	Mean Low Water Springs
MNCR	Marine Nature Conservation Review
MPA	Marine Protected Area
MSL	Mean Sea Level
MS-LOT	Marine Scotland Licensing Operations Team
nm	Nautical Mile
PMF	Priority Marine Feature
Ramsar	Wetlands of international importance, designated under The Convention on Wetlands (Ramsar, Iran, 1971)
SAC	Special Areas of Conservation
SACFOR	Super-abundant, Abundant, Common, Frequent, Occasional, Rare (and less than rare)
SNH	Scottish Natural Heritage
SPA	Special Protection Area
SSSI	Site of Special Scientific Interest
UKHO	United Kingdom Hydrographic Office
WFD	Water Framework Directive

Cardinal points/directions are used unless otherwise stated.

SI units are used unless otherwise stated.

Appendices



Innovative Thinking - Sustainable Solutions

A Drop-Down Camera Survey Report

A.1 Introduction

This survey report reviews the baseline ecological characteristics of the subtidal habitats in the vicinity of the Hunterston B Power Station cooling water intake. The specific objectives of the survey were to achieve the following:

- Characterise the density and distribution of kelp species within the survey area (proposed seaweed reduction area);
- Produce a broad-scale map of subtidal habitats; and
- Confirm the presence and distribution of any protected or nationally scarce biotopes such as maerl beds, horse mussel beds or seagrass beds.

A.2 Survey Methodology

The drop-down camera survey was undertaken on 31 May 2017 using the commercial fishing vessel *Eilidh Anne*. The survey approach was based on the procedures identified in the Marine Monitoring Handbook, Procedural Guidance No 3-5 (Holt and Sanderson, 2001) and guidance prepared by the Northeast Atlantic Marine Biological Analytical Quality Control Scheme (NMBAQC) (Hitchin et al. 2015).

The survey used a drop-down underwater camera with a cable to the vessel to allow video footage to be viewed in real time. In addition, a Go Pro action camera (with red filter) was also used which was collecting both video and still images. The cameras were mounted on a mobile frame unit (Image A.1).

For shallow infralittoral areas with the highest potential kelp densities (depths of approximately 4-6 m), the frame was fixed below the hull of the boat and towed slowly at approximately 1-2 knots along two transect lines (T1 and T2). An error with the vessels data communication tracking system occurred on the day of the survey; therefore, the actual position of the vessel along each of the transect lines was derived from positions inputted into a handheld Garmin Global Positioning System (GPS) (see Image A.2 for site locations).



Image A.1. Mobile frame used for drop down video survey




Image A.2. Transect and drop-down video survey points

For deeper areas (greater than approximately 8 m depth), the frame was dropped at fixed intervals, towed for a short distance before being retrieved. In total, 79 stations were sampled using this technique.

At each location, the density and coverage of kelp was recorded based on the Marine Nature Conservation Review (MNCR) SACFOR abundance scale (Table A.1). Biotores were also assigned based on the Marine Habitat Classification for Britain and Ireland (MHCBI) 04.05 to biotope class levels 4 or 5 (Conner et al. 2004).

Table A.1. SACFOR Abundance Scale

Code	Description	Coverage
S	Super-abundant	>80%
A	Abundant	40-79%
C	Common	20-39%
F	Frequent	10-19%
O	Occasional	5-9%
R	Rare	1-5% or density
L	Less than rare (indicated by extrapolation)	<1% or density

A.3 Results

In total, over six hours of video footage was collected from the survey and analysed. Annex A.1 presents a summary of the survey results. This includes details of the depth, kelp coverage (based on the SACFOR scale), habitat type and species observed, as well as still images from the video at representative locations along the transects and also for each of the drop-down locations.

To further summarise the key findings from this work, the survey area was divided into three broad areas based on the depth profile. These areas are as follows:

- Shallow infralittoral (approx. 4-7 m depth);
- Mid infralittoral (approx. 8-15 m depth); and
- Lower infralittoral (depths deeper than approx. 15 m).

The broad habitats recorded at each of these depths are described in more detail below. It should be noted that no protected or nationally scarce habitats were recorded in the survey.

A.3.1 Shallow Infralittoral

The habitat within the shallow infralittoral to the north of the Hunterston B cooling water intake was characterised by mixed sediment (predominantly sand with gravel and cobbles along with boulders and shell debris). Within this area, sugar kelp *Saccharina latissima* (*Laminaria saccharina*) coverage was very dense (Image A.3). Occasional oarweed *Laminaria digitata* was also present attached to larger boulders. Based on the SACFOR abundance scale, kelp density generally varied from abundant to super-abundant (coverage >40%) in this area. Kelp fronds were approximately 1 to 3 m in length (Image A.4).




Image A.3. Dense kelp coverage within the infralittoral north of the cooling water intake



Image A.4. Example of *Saccharina latissima* frond on deck of survey vessel

Red seaweeds (such as *Plocamium cartilagineum*, *Polysiphonia elongata* and *Lomentaria clavellosa*) and brown seaweeds (including dead man's rope *Chorda filum*) were frequently observed in between kelp patches and under the kelp canopy. Epifaunal species commonly recorded included spiny starfish *Marthasterias glacialis* and edible sea urchin *Echinus esculentus*. The invasive wireweed *Sargassum muticum* was also recorded on several occasions. Mobile species recorded included the edible crab *Cancer pagurus*, shore crab *Carcinus maenas*, small-spotted catshark *Scyliorhinus canicula* and various wrasse species.

This habitat is most appropriately assigned to SS.SMp.KSwSS.LsacR (*Laminaria saccharina* and red seaweeds on infralittoral sediments) and IR.LIR.K.Lsac.Ldig (*Laminaria saccharina* and *Laminaria digitata* on sheltered sublittoral fringe rock).

Sugar kelp *S. latissima* coverage was generally less dense and patchier to the south of the Hunterston B cooling water intake (Image A.5). Kelp density generally varied from frequent to occasional in this area based on the SACFOR abundance scale (coverage >20%). The sediment generally had less cobbles and boulders compared with the area north of the Hunterston B cooling water intake, although several boulder fields were observed with denser kelp coverage evident. Spiny starfish *M. glacialis* and edible sea urchin *E. esculentus* were frequently recorded, with the common starfish *Asteria rubens* recorded occasionally. On several occasions, dead man's fingers *Alcyonium digitatum* were recorded attached to larger cobbles. Towards the south of the towed transect in this area, kelp coverage was particularly sparse with the area characterised predominately by red and brown algae species and spiny starfish.

The shallow infralittoral habitat to the south of the Hunterston B cooling water intake is most approximately assigned to SS.SMp.KSwSS.LsacR.S (*Laminaria saccharina* and filamentous red algae on infralittoral sand).




Image A.5. Spiny starfish *Marthasterias glacialis* and sugar kelp *Saccharina latissima* frond

A.3.2 Mid Infralittoral

At depths of approximately 8-15 m (mid infralittoral), coverage of sugar kelp *S. latissima* was generally very sparse with only occasional fronds recorded (coverage on the SACFOR abundance scale of <5%, i.e. classified as rare). The substrate consisted of silty sand with shell debris. Brown algae along with the sea beard *Nemertesia antennina*, spiny starfish *M. glacialis* and common starfish *A. rubens* were frequently recorded (Image A.6). Dead man's fingers *Alcyonium digitatum* and the sponge *Suberites carnosus* were also occasionally observed. This habitat has been assigned to SS.SSa.IMuSa (Infralittoral muddy sand).

A.3.3 Lower Infralittoral

No kelp was evident at the deeper depths surveyed, with the habitat consisting of sandy silt (SS.SMu.ISaMu: Infralittoral sandy mud, see Image A.7).



Image A.6. Spiny starfish *Marthasterias glacialis* and common starfish *Asteria rubens* with shell debris



Image A.7. Sandy mud habitat recorded at deeper depths sampled during survey

A.3.4 Other Ecological Observations

A common seal *Phoca vitulina* was observed hauled out at low water on a flat rock below the Hunterston B Power Station (in the vicinity of the cooling water outfall). A solitary short-beaked common dolphin *Delphinus delphis*, known locally as 'Kylie', was also observed while on the return journey to Largs Marina. In addition, large numbers of moon jellyfish *Aurelia aurita*, lion's mane jellyfish *Cyanea capillata*, comb jellies and sea gooseberrys were observed.

A.4 References

Connor, D.W., Allen, J.H., Golding, N., Howell, K.L., Lieberknecht, L.M., Northen, K.O. and Reker, J.B. 2004. The Marine Habitat Classification for Britain and Ireland Version 04.05. In: Joint Nature Conservation Committee (JNCC). 2015. The Marine Habitat Classification for Britain and Ireland Version 15.03. Available online at: jncc.defra.gov.uk/MarineHabitatClassification (Accessed June 2017).

Hitchin, R., Turner, J.A. and Verling, E. 2015. NMBAQC/JNCC Epibiota remote monitoring from digital imagery: Operational guidelines. 25pp, July 2015

Holt, R. and Sanderson, B. 2001. Procedural Guideline No. 3-5 - Procedural Guideline No. 3-5. Identifying biotopes using video recordings. In: Davies J., Baxter J., Bradley M., Connor D., Khan J., Murray E., Sanderson W., Turnbull C. and Vincent M. 2001. Marine Monitoring Handbook, 405 pp.



A.5 Abbreviations/Acronyms

GPS	Global Positioning System
MHCBI	Marine Habitat Classification for Britain and Ireland
MNCR	Marine Nature Conservation Review
NMBAQC	Northeast Atlantic Marine Biological Analytical Quality Control Scheme (NMBAQC)
SACFOR	Abundance scale: Super-abundant (S), Abundant (A), Common (C), Frequent (F), Occasional (O), Rare (R), Less than rare (L)

Cardinal points/directions are used unless otherwise stated.

SI units are used unless otherwise stated.

Annex A.1 Survey Results

Site	Approx. Depth	Kelp Coverage ¹	Biotope	Survey Notes	Drop-Down Video Still Frame
T1-a		S-A	SS.SMp.KSwSS.LsacR (<i>Laminaria saccharina</i> and red seaweeds on infralittoral sediments)	Sand with gravel and cobbles. Dense coverage of sugar kelp <i>Saccharina latissima</i> and very occasional oarweed <i>Laminaria digitata</i> . Red seaweeds and brown seaweeds (including dead man's rope <i>Chorda filum</i> . Spiny starfish <i>Marthasterias glacialis</i> and edible sea urchin <i>Echinus esculentus</i> observed frequently.	
T1-a		S-A	SS.SMp.KSwSS.LsacR (<i>Laminaria saccharina</i> and red seaweeds on infralittoral sediments)	Sand with gravel and cobbles. Dense coverage of sugar kelp <i>Saccharina latissima</i> and very occasional oarweed <i>Laminaria digitata</i> . Red seaweeds and brown seaweeds (including dead man's rope <i>Chorda filum</i> . Spiny starfish <i>Marthasterias glacialis</i> and edible sea urchin <i>Echinus esculentus</i> observed frequently.	
T1-b		S-A	SS.SMp.KSwSS.LsacR (<i>Laminaria saccharina</i> and red seaweeds on infralittoral sediments)	Sand with gravel and cobbles. Dense coverage of sugar kelp. Brown and red seaweeds. Spiny starfish, common sea urchin.	